

ORNSTEIN-UHLENBECK PROCESSES WITH SINGULAR DRIFTS: INTEGRAL ESTIMATES AND GIRSANOV DENSITIES

MARIA GORDINA*, MICHAEL RÖCKNER‡, AND ALEXANDER TEPLYAEV†

ABSTRACT. We consider a perturbation of a Hilbert space-valued Ornstein–Uhlenbeck process by a class of singular nonlinear non-autonomous maximal monotone time-dependent drifts. The only further assumption on the drift is that it is bounded on balls in the Hilbert space uniformly in time. First we introduce a new notion of generalized solutions for such equations which we call pseudo-weak solutions and prove that they always exist and obtain some pathwise estimates in terms of the data of the equation. Then we prove that their laws are absolutely continuous with respect to the law of the original Ornstein–Uhlenbeck process. In particular, we show that pseudo-weak solutions always have continuous sample paths. In addition, we obtain integrability estimates of the associated Girsanov densities. Some of our results concern non-random equations as well, while probabilistic results are new even in finite-dimensional autonomous situations.

CONTENTS

1. Introduction	2
Acknowledgements	3
2. Setting and main results	3
2.1. Setting and assumptions	3
2.2. Pseudo-weak solutions and their properties	4
3. Preliminaries: Pseudo-weak convergence and Yosida approximations	6
3.1. Pseudo-weak convergence	6
3.2. Yosida approximations to A	9
3.3. Yosida approximations to F	10
4. Almost sure φ -type estimates of solutions X_t	11
4.1. Properties of function space \mathcal{M}	11
4.2. Proof of Theorem 2.8	14
4.3. Further <i>a priori</i> pathwise estimates of X_t and proof of Theorem 2.9	16
5. Uniform integrability of Girsanov densities	16
5.1. Stopping times and Girsanov transforms	16
5.2. Estimates of the Girsanov densities and proof of Theorem 2.10	18
5.3. Quantitative estimates and examples	21
References	23

1991 *Mathematics Subject Classification*. Primary 60H10; Secondary 35R15, 60H15, 47D07, 47N30.

Key words and phrases. Ornstein-Uhlenbeck process, singular perturbation, nonlinear infinite-dimensional stochastic differential equations, non-Lipschitz monotone coefficients, Girsanov theorem.

*Research was supported in part by NSF Grants DMS-1405169, DMS-1712427, and a Simons Fellowship.

‡ Research was supported in part by the German Science Foundation (DFG) through CRC 1283.

† Research was supported in part by NSF Grant DMS-1613025.

1. INTRODUCTION

Suppose H is a real separable Hilbert space with an inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $|\cdot|_H$. The aim of this paper is to study solutions to the following H -valued stochastic differential equation

$$(1.1) \quad dX_t = (AX_t + F(t, X_t)) dt + \sigma dW_t, \quad X_0 = x \in H, t \geq 0.$$

Here W_t is a cylindrical Wiener process in H on some filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$ satisfying the usual conditions of right continuity and \mathbb{P} -completeness. For the precise setting we refer to Section 2.1.

We consider Equation (1.1) without the standard assumption on $F(t, \cdot)$ being locally Lipschitz continuous. The motivation for our study includes a better understanding of equations such as (1.1) with time-dependent drifts of not necessarily polynomial growth.

Equation (1.1) can be viewed as a nonlinear non-autonomous perturbation of the stochastic differential equation corresponding to an Ornstein-Uhlenbeck process. In fact, it is a long-standing open problem to find optimal or nearly optimal conditions on F such that (1.1) has a solution under the usual assumption that A generates a C_0 -semigroup on H (see e.g. [15, 18] and the references therein). If F is maximal monotone and single-valued with $D_F = H$, then we can equivalently rewrite (1.1) as the random equation

$$(1.2) \quad dZ_t = (AZ_t + F(t, Z_t + W_{0,A,\sigma}(t))) dt, \quad Z_0 = x,$$

where $Z_t = X_t - W_{0,A,\sigma}(t)$ and the Ornstein-Uhlenbeck process $W_{0,A,\sigma}$ solves (1.1) for $F \equiv 0$, $x = 0$. Moreover, in this case one can easily obtain a unique solution by classical results due to F. Browder, Kato, Komura and Rockafellar in [11, 28, 29, 40]. However, the assumption $D_F = H$ excludes many interesting examples [15] and [18, Section 7.2], and therefore we include the case $D_F \subsetneq H$, and also allow F to be multi-valued, see Assumptions 2.3 and 2.4 below. The first main result of our paper is that under natural assumptions on F , namely, Assumptions 2.3 and 2.4, (1.1) always has a solution in a generalized sense. Namely, we introduce **pseudo-weak solutions** in Definition 2.5 and discuss them in detail in Section 3.1.

The main results of the paper include a proof of existence of pseudo-weak solutions, pathwise *a priori* estimates of these solutions in Section 4, absolute continuity of the law of these solutions with respect to the law of the Ornstein-Uhlenbeck process, and finally integral estimates of the corresponding Radon-Nikodym densities in Section 5. Our approach can be interpreted as an extension of the classical use of Girsanov transformation to find a solution for a stochastic differential equation with a nonzero (but at most linearly growing) drift. The main idea behind results such as Theorem 2.8 is that we can estimate $\varphi(|X_t|_H^2)$, where X_t is a solution to Equation 1.1 and φ is a suitable function. The function φ can be chosen by looking at the behavior of the nonlinearity F at infinity, so the estimate only depends on the growth of F at infinity, not on the nonlinearity itself.

Some of our results on the Girsanov transform are closely related to the infinite-dimensional estimates by D. Gatarek and B. Gołdys in [24, 25]. They considered equations in Banach spaces, while we restrict our consideration to Hilbert space though for non-autonomous perturbations. In the future work we plan to extend our techniques to the reaction-diffusion equation in a Banach space. Our estimates of solutions and Girsanov densities are new even in finite dimensions, for example, compared to the ones due to N. V. Krylov in [30, 32] and [31, Chapter IV, §3].

We would like to comment on some of the previous results both in terms of the assumptions we make and the techniques we use. We describe the setting in Section 2.1 in detail, including the assumptions on the coefficients of the non-autonomous equation (1.1). The approach we use does not rely on an invariant measure, which is not available for non-autonomous equations, and therefore we do not use typical assumptions such as finite moments of the invariant measure and integrability properties of the nonlinear drift with respect to this measure. The paper consists of three major parts which are intertwined: in Section 2.2 we introduce a notion of pseudo-weak solutions to (1.1), and prove their existence in Section 4.3. We use monotonicity of the coefficients of the equation to prove *a priori* pathwise bounds in Theorem 2.8. In general one expects that our assumptions might imply weak uniqueness, by appealing to Gronwall's lemma, but this seems out of reach for now in a general setting such as ours. We refer to [8, 15] for a discussion of when and how martingale solutions to (1.1) can be constructed, and for more details on such solutions.

We also would like to mention several connections of our results to the cases when F in (1.1) does not depend on t . In this situation one can use our results to prove smoothness results for an invariant measure, closability of the corresponding Dirichlet form etc. Note that in the current paper we do not address the question under which assumptions an invariant measure exists. Suppose there is an invariant measure as described in [15], then one can use the Girsanov transform in Theorem 2.10 to show formally that the invariant measure is quasi-invariant under certain linear shifts. This leads to a possibility of using [3, Theorem 2.2] and [1, Theorem 1.3] to prove closability of the Dirichlet form. The main ingredient here is the lower semicontinuity of the Radon-Nikodym density as described in [2, p.122] among other references. Finally, there are other recent approaches to quasi-invariance of semigroups in infinite dimensions which rely on functional inequalities [6, 7, 20, 21, 26, 27, 35, 42], and these methods are not applicable to singular perturbations considered in the current paper.

Acknowledgements. The authors thank G. Da Prato, M. Hairer, M. Hinz and N. Krylov for helpful discussions and suggestions. The authors are grateful to anonymous referees for corrections and suggested improvements to the paper.

2. SETTING AND MAIN RESULTS

2.1. Setting and assumptions. Let H be a real separable Hilbert space with an inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $|\cdot|_H$. We denote by $B(H)$ the space of bounded linear operators equipped with the operator norm $\|\cdot\|$. The Hilbert-Schmidt norm is denoted by $\|\cdot\|_{HS}$. We suppose that the coefficients A , F and B in Equation (1.1) satisfy the following assumptions.

Assumption 2.1. *The operator (A, D_A) generates a C_0 -semigroup on H denoted by e^{tA} , $t \geq 0$. We assume that there is $\beta > 0$ such that for all $x \in D_A$*

$$\langle Ax, x \rangle \leq -\beta|x|_H^2.$$

Note that Assumption 2.1 implies that A is m -dissipative on H .

Assumption 2.2. *Both σ and σ^{-1} are in $B(H)$ with σ being self-adjoint and positive such that for some $\varepsilon > 0$*

$$\int_0^T t^{-\varepsilon} \|e^{tA}\sigma\|_{HS}^2 dt < \infty, \text{ for all } T > 0.$$

Recall that under Assumption 2.2 the Ornstein-Uhlenbeck process

$$(2.1) \quad W_{x,A,\sigma}(t) := e^{tA}x + \int_0^t e^{(t-s)A}\sigma dW_s, \quad t \geq 0,$$

is pathwise continuous by [14, Proposition 2.3] which is based on the technique described in [13].

Assumption 2.3. Denote by 2^H the power set of the Hilbert space H . Let $F(t, \cdot) : [0, \infty) \times D_F \rightarrow 2^H$ be a family of maps such that D_F is a non-empty Borel set in H , and $dt \otimes \mathbb{P}$ -almost surely the Ornstein-Uhlenbeck process $W_{x,A,\sigma} \in D_F$ for all $x \in D_F$. Furthermore, $F(t, \cdot)$ is an m -dissipative map, that is, for any $x_1, x_2 \in D_F$

$$\langle y_1 - y_2, x_1 - x_2 \rangle \leq 0, \quad \text{for any } y_1 \in F(t, x_1), y_2 \in F(t, x_2), t \in [0, \infty)$$

and for any $\alpha > 0$ and $t \in [0, \infty)$

$$\text{Range}(\alpha I - F(t, \cdot)) := \{\alpha x - y : y \in F(t, x), x \in D_F\} = H.$$

We refer to [4, Section II.3] and [5, Chapter 3] for basic facts about dissipative maps, as well as to the exposition in [45]. In particular, it is known that in a Hilbert space a map is m -dissipative if and only if it is maximal dissipative, that is, it has no proper dissipative extensions. By [4, Proposition 3.5(iv), Chapter II] for any $(t, x) \in [0, \infty) \times D_F$, the set $F(t, x)$ is non-empty, closed and convex, and so we can consider the well-defined single-valued map

$$(2.2) \quad F_0(t, x) := \{y \in F(t, x) : |y|_H = \inf\{|z|_H, z \in F(t, x)\}\} \quad \text{for any } x \in D_F.$$

This definition is the same as in [15] except that we allow dependence on time. Using the Yosida approximation to F described in Section 3 we see that function $F_0(t, x)$, usually called *the minimal section of the maximal monotone operator F* , is Borel measurable. Our next assumption is similar to the ones introduced in [17, 25].

Assumption 2.4. $F_0(t, \cdot)$ is uniformly bounded in $t \in [0, \infty)$ on balls in H , that is,

$$(2.3) \quad a(r) := \sup_{x \in B_r \cap D_F} \sup_{t \in [0, \infty)} |F_0(t, x)|_H < \infty,$$

where $B_r := \{x \in H : |x|_H < r\}$, $r > 0$.

We define $a(0) := 0$. The function a is non-decreasing and left-continuous, and therefore Borel measurable.

We are mostly interested in the case when $\lim_{u \rightarrow \infty} a(u) = \infty$. Assumption (2.4) simply means that $F_0(t, x)$ is bounded on balls in its domain of definition in x , uniformly in t . In other words, we assume that function $F_0(\cdot, \cdot)$ is locally bounded in the space variable uniformly in time.

2.2. Pseudo-weak solutions and their properties. Throughout this paper we assume that Assumptions 2.1, 2.2, 2.3 and 2.4 hold. The first step in defining pseudo-weak solutions to Equation (1.1) requires suitable approximations to F . We use the Yosida approximation $F_\alpha, \alpha > 0$ described in Section 3.3 below.

By $Z_{\alpha,t}^x$ we denote the continuous H -valued process which is a mild solution to the family of regularized random ordinary differential equations

$$(2.4) \quad dZ_{\alpha,t}^x = (AZ_{\alpha,t} + F_\alpha(t, Z_{\alpha,t}^x + W_{0,A,\sigma}(t))) dt, \quad Z_{\alpha,0}^x = x,$$

where $W_{0,A,\sigma}$ is the pathwise continuous Ornstein-Uhlenbeck process defined by (2.1) with $x = 0$. One can use [39, Chapter 6, Theorem 1.2, page 184] to justify the existence of mild solutions to (2.4). We note that technically speaking [39] assumes that F_α is continuous in time, but it is clear that this assumption is not essential, and it is enough to assume joint measurability in time and space, and Lipschitz continuity in space, with the Lipschitz constant uniform in time, which holds for F_α as we explain in Section 3.3.

The stochastic differential equation

$$(2.5) \quad \begin{aligned} dX_t &= (AX_t + F_\alpha(t, X_t)) dt + \sigma dW_t, \\ X_0 &= x \in H \end{aligned}$$

has a mild solution $X_{\alpha,t}^x = X_\alpha(t, x)$, $t \geq 0$, with \mathbb{P} -a.s. continuous sample paths. Even though we have dependence on α in this equation, we prove that solutions $X_{\alpha,t}^x$ satisfy bounds (2.6), which are uniform in α . In addition, it is clear that $Z_{\alpha,t}^x$ is a mild solution to the random ordinary differential equation (2.4) if and only if

$$X_{\alpha,t}^x := Z_{\alpha,t}^x + W_{0,A,\sigma}(t)$$

is a mild solution to (2.5).

Before proceeding to the notion of pseudo-weak solutions, we would like to comment on the intuition behind it. First we introduce pseudo-weak limits to deal with non-metrized topology. One of the consequences of this definition is that the convergence is governed by a function ψ , and therefore the limit might be different for different choices of ψ as we discuss later. In what follows, unless stated otherwise, a pseudo-weak limit means a ψ -pseudo-weak limit, in the sense of Definition 3.2 and Remark 3.3 below.

Definition 2.5. An adapted H -valued process X^x is a *pseudo-weak solution* to (1.1) if it is a pseudo-weak limit point in $L^2([0, \infty) \times \Omega, dt \times \mathbb{P}; H)$ of the approximating processes X_α^x defined by (2.5).

Remark 2.6. Obviously, such pseudo-weak limit points are automatically adapted. More surprisingly, Theorem 2.10 implies that they are also continuous \mathbb{P} -a.s. in H .

The main results of our paper are summarized in the following three theorems. We start with pathwise *a priori* estimates. For this purpose we introduce the function space \mathcal{M} as follows.

Definition 2.7. Let \mathcal{M} denote the space of continuous functions $\varphi : [0, \infty) \rightarrow (0, \infty)$ such that

- (1) φ is a strictly increasing convex function which is C^2 on $(0, \infty)$;
- (2) the limit $\frac{u\varphi'(u)}{\varphi(u)} \xrightarrow{u \rightarrow \infty} L_\varphi$ exists, and $L_\varphi \in [1, \infty]$.

For properties and examples of functions in \mathcal{M} we refer to Section 4.1.

Theorem 2.8 (Uniform pathwise *a priori* φ -estimates). *Under Assumptions 2.1, 2.2, 2.3, 2.4, for every $\varphi \in \mathcal{M}$ we have the following estimates for a pseudo-weak solution X_t^x to Equation (1.1)*

$$(2.6) \quad \varphi(|X_t^x|_H^2) \leq \frac{e^{-\beta t}}{2} \varphi(4|x|_H^2) + \frac{1}{2} K_\varphi(t) + \frac{\beta t}{2} K_{\varphi,\beta,a}(t) < \infty \text{ a.s.}$$

Here $K_\varphi(t)$ and $K_{\varphi,\beta,a}(t)$ are random functions defined in Notation 4.6 below. These functions only depend on β, σ, A and a .

Theorem 2.9 (Pseudo-weak solutions). *Under Assumptions 2.1, 2.2, 2.3, 2.4, there exists a pseudo-weak solution $\{X_t^x\}_{t \geq 0}$ to Equation (1.1), i.e.*

$$(2.7) \quad X_t^x := Z_t^x + W_{0,A,\sigma}(t) \quad t \geq 0,$$

where the process Z_t^x is a pseudo-weak limit point of $Z_{\alpha,t}^x$, as $\alpha \rightarrow 0$, and $Z_{\alpha,t}^x$ is a solution to Equation (2.4). Moreover, $(dt \times \mathbb{P})$ -a.s. we have the following estimate

$$(2.8) \quad |X_t^x|_H \leq |x|_H e^{-\beta t} + \int_0^t e^{-\beta(t-s)} a(|W_{0,A,\sigma}(s)|_H) ds + |W_{0,A,\sigma}(t)|_H.$$

In the next theorem we prove a Girsanov-type result with respect to the law of the Ornstein–Uhlenbeck process $W_{x,A,\sigma}$ defined by (2.1). As we mentioned earlier, we can view this result as an analogue of using a Girsanov transformation to find a solution to stochastic differential equation where the reference process is the Ornstein–Uhlenbeck process.

Theorem 2.10. *Suppose Assumptions 2.1, 2.2, 2.3, 2.4 hold. Then on any finite time interval $[0, T]$ and for any $x \in D_F$ the law of a pseudo-weak solution X_t^x to Equation (1.1) is absolutely continuous with respect to the law of $W_{x,A,\sigma}$ on $L^2([0, T], dt; H)$. In addition, the solution X_t^x has \mathbb{P} -a.s. continuous sample paths in H .*

Remark 2.11. One can expect that the corresponding Radon-Nikodym densities $\{\rho^x\}_{x \in B_r}$ are uniformly integrable for any ball B_r , $r > 0$. Note that uniform integrability of the Radon-Nikodym densities $\{\rho^x\}_{x \in B_r}$ in Theorem 2.10 holds if and only if for any $r > 0$ there exists an increasing unbounded function $\Psi : [0, \infty) \rightarrow [0, \infty)$ such that $\sup_{x \in B_r} \mathbb{E} \rho^x \Psi(\rho^x) < \infty$. In our paper we prove a weaker estimate

$$(2.9) \quad \mathbb{E} \rho^x \Psi(\rho^x) < \infty.$$

In Subsection 5.3 we give quantitative estimates of $\Psi(\cdot)$ in terms of the function $a(\cdot)$ and the tail probability estimates of the Ornstein-Uhlenbeck process $W_{x,A,\sigma}$ under natural additional assumptions.

We stress that Theorem 2.10 is to the best of our knowledge new, even if $H = \mathbb{R}^d$.

We prove Theorem 2.8 and Theorem 2.9 in Section 4, where we provide more detailed statements as well. These results are illustrated by Examples 4.2, 4.3, 4.4. Theorem 2.10 is proved in Section 5.2 and an example of how to construct Ψ is given in Example 5.4. Note that Theorem 2.10 addresses the absolute continuity of the laws which is a long-standing question that has been implicitly stated in a number of publications such as [43, 44].

3. PRELIMINARIES: PSEUDO-WEAK CONVERGENCE AND YOSIDA APPROXIMATIONS

3.1. Pseudo-weak convergence. Let (S, \mathcal{F}, μ) be a σ -finite measure space, then for any $A \in \mathcal{F}$ we set

$$\mu_A := \mathbb{1}_A \mu.$$

Let H be a separable real Hilbert space with an inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $|\cdot|_H$.

Notation 3.1. We need several spaces of H -valued functions on the measure space (S, \mathcal{F}, μ) . By $L^0(S, \mu; H)$ we denote the space of equivalence classes of \mathcal{F} -measurable function on (S, \mathcal{F}, μ) defined up to sets of μ -measure zero and equipped with the topology of convergence

in measure. By $L^2(S, \mu; H)$ we denote the space of H -valued square-integrable functions on S .

In what follows we use $\xrightarrow[n \rightarrow \infty]$ for the weak convergence in Banach spaces.

Definition 3.2. Suppose $f, f_n \in L^0(S, \mu; H)$, $n \in \mathbb{N}$. We say that $\{f_n\}_{n=1}^\infty$ converges *pseudo-weakly* to f , denoted by

$$f_n \xrightarrow[n \rightarrow \infty]{\psi} f,$$

if

$$(3.1) \quad \psi(f_n) \xrightarrow[n \rightarrow \infty]{} \psi(f) \text{ in } L^2(S, \mu_A; H) \text{ for any } A \in \mathcal{F} \text{ with } \mu(A) < \infty$$

and for some $\psi : H \rightarrow H$ defined by

$$(3.2) \quad \psi(h) := \begin{cases} \frac{h}{|h|_H} \psi_0(|h|_H), & \text{if } h \neq 0, \\ 0, & \text{if } h = 0, \end{cases}$$

where $\psi_0 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a strictly increasing continuous function such that $\psi_0(0) = 0$. In this case we say that f is a ψ -pseudo-weak limit of the sequence $\{f_n\}_{n=1}^\infty$.

Remark 3.3. The definition of ψ -pseudo-weak limits depends on the choice of function ψ which we usually fix. Typical examples for such a function ψ are $\psi(h) = h$ or

$$(3.3) \quad \psi(h) = \frac{h}{1 + |h|_H}, h \in H.$$

The latter choice corresponds to $\psi_0(r) = \frac{r}{1+r}$ for $r \in \mathbb{R}_+$. For us the most interesting case is when ψ is bounded, and most of the time we use ψ defined by (3.3), in which case we say *pseudo-weak* convergence dropping the dependence on ψ .

Remark 3.4. The standard weak convergence coincides with ψ -pseudo-weak convergence for $\psi(h) = h$. For a space such as $H = L^2(\mathbb{R}^d)$, the converse can be proved as well: if the ψ -pseudo-weak convergence coincides with the usual weak convergence then $\psi(h) = h$ up to a multiplicative constant. We do not intend to study this question in detail in this paper, and only mention it in order to provide a better intuition for this notion of convergence.

Proposition 3.5. Suppose $f, f_n \in L^0(S, \mu; H)$. Then for any bounded $\psi : H \rightarrow H$ we have that $f_n \xrightarrow[n \rightarrow \infty]{\psi} f$ if and only if

$$(3.4) \quad \int_A \langle \psi(f_n) - \psi(f), h \rangle d\mu \xrightarrow[n \rightarrow \infty]{} 0$$

for any $h \in H$ and any $A \in \mathcal{F}$ with $\mu(A) < \infty$.

Remark 3.6. Observe that for a fixed ψ the pseudo-weak limit is *unique*, that is, if

$$f_n \xrightarrow[n \rightarrow \infty]{\psi} f,$$

$$f_n \xrightarrow[n \rightarrow \infty]{\psi} g,$$

then $f = g$ μ -a.e.

Remark 3.7. Note that, the topology of $L^0(S, \mu; H)$ defined by convergence in measure

$$\lim_{n \rightarrow \infty} \mu(\{|f_n - f|_H > \varepsilon\} \cap A) = 0 \quad \text{for all } \varepsilon > 0, A \in \mathcal{F}, \mu(A) < \infty$$

implies pseudo-weak convergence, but these two types of convergence are not equivalent in general.

The following assertion is an easy consequence of the Banach-Saks Theorem applied to the Hilbert space $L^2(S, \mu; H)$ or, more elementarily, of Fatou's Lemma.

Proposition 3.8. *Suppose $f, f_n \in L^2(S, \mu; H)$, $n \in \mathbb{N}$, and*

$$(3.5) \quad f_n \xrightarrow[n \rightarrow \infty]{} f,$$

then

$$|f|_H \leq \limsup_{n \rightarrow \infty} |f_n|_H \quad \mu\text{-a.e.}$$

Corollary 3.9. *Let $f, f_n \in L^0(S, \mu; H)$, $n \in \mathbb{N}$, such that*

$$f_n \xrightarrow[n \rightarrow \infty]{\psi} f.$$

Then

$$|f|_H \leq \limsup_{n \rightarrow \infty} |f_n|_H \quad \mu\text{-a.e.}$$

Proof. Let $A \in \mathcal{F}$, $\mu(A) < \infty$ and ψ as in Definition 3.2. Then by Proposition 3.8 applied to μ_A instead of μ we have that on the set

$$\left\{ f \neq 0, \limsup_{n \rightarrow \infty} |f_n|_H < \infty \right\}$$

we have

$$\begin{aligned} 0 < \psi_0(|f|_H) &= |\psi(f)|_H \leq \limsup_{n \rightarrow \infty} |\psi(f_n)|_H \\ &= \limsup_{n \rightarrow \infty} \psi_0(|f_n|_H) \leq \psi_0\left(\limsup_{n \rightarrow \infty} |f_n|_H\right) \quad \mu_A - \text{a.e.} \end{aligned}$$

Applying the inverse of ψ_0 to both sides of this inequality and using the fact that μ is σ -finite proves the desired result. \square

Proposition 3.10. *Let ψ in Definition 3.2 be bounded. If $f_n \in L^0(S, \mu; H)$, $n \in \mathbb{N}$, are such that*

$$\sup_{n \in \mathbb{N}} |f_n|_H < \infty \quad \mu - \text{a.e.},$$

then there exists $f \in L^0(S, \mu; H)$ such that for some subsequence $\{n_k\}_{k \in \mathbb{N}}$

$$f_{n_k} \xrightarrow[k \rightarrow \infty]{\psi} f.$$

Proof. Let $B_R(0)$ denote the open ball in H with center 0 and radius $0 < R < \infty$. Define $\psi^{-1} : B_{|\psi|_\infty} \rightarrow H$ by

$$\psi^{-1}(h) := \begin{cases} \frac{h}{|h|_H} \psi_0^{-1}(|h|_H), & \text{if } h \neq 0, \\ 0, & \text{if } h = 0, \end{cases}$$

where ψ_0^{-1} is the inverse function of ψ_0 . It is easy to see that ψ^{-1} is indeed the inverse map of ψ with ψ defined by (3.2). Now let $A \in \mathcal{F}$, $\mu(A) < \infty$, and

$$V_n := \psi(f_n), \quad n \in \mathbb{N}.$$

Then $\{V_n\}_{n \in \mathbb{N}}$ is bounded in $L^2(S, \mu_A; H)$. Hence there exists a $V_A \in L^2(S, \mu_A; H)$ such that for some subsequence $\{n_k\}_{k \in \mathbb{N}}$

$$V_{n_k} \xrightarrow{m \rightarrow \infty} V_A$$

in $L^2(S, \mu_A; H)$. Since μ is σ -finite, we can choose a sequence of subsets A_l of finite μ -measure such that $\cup A_l = S$, and by a diagonal argument we can construct $V \in L^2(S, \mu; H)$ and a subsequence, again indexed by $n_k, k \in \mathbb{N}$ such that

$$V_{n_k} \xrightarrow{k \rightarrow \infty} V.$$

By Proposition 3.8 we have that

$$|V|_H \leq \limsup_{k \rightarrow \infty} |V_{n_k}|_H = \limsup_{k \rightarrow \infty} \psi_0(|f_{n_k}|_H) \leq \psi_0(\sup_{n \in \mathbb{N}} |f_n|_H) \quad \mu\text{-a.e.}$$

Therefore $V \in B_{|\psi_0|_\infty}(0)$ and

$$f := \psi^{-1}(V)$$

is well-defined. By definition of the ψ -pseudo-weak convergence

$$f_{n_k} \xrightarrow[k \rightarrow \infty]{\psi} f.$$

□

3.2. Yosida approximations to A . We need the Yosida approximations A_α to A for small α , in particular, we will use the fact that such A_α satisfy Assumption 2.1 with a change of β as in Proposition 3.11. Surprisingly, it is not easy to find a reference to this fact, so again we include it for completeness.

We start by recalling some standard facts about C_0 -semigroups and their generators, most of this goes back to Hille and Yosida. We refer to [22, Chapter II] for most of the material below. Let $\rho(A)$ be the resolvent set, then the resolvent of A is defined as

$$\begin{aligned} R_\lambda(A) &:= (\lambda I - A)^{-1}, \quad \lambda \in \rho(A) \in B(H), \\ R_\lambda(A) &: H \longrightarrow D_A. \end{aligned}$$

Recall that for $\lambda > 0$ we have $\|R_\lambda(A)\| \leq 1/\lambda$. In addition,

$$(3.6) \quad \lambda R_\lambda(A) x \xrightarrow{\lambda \rightarrow \infty} x, \quad x \in H.$$

Note that $AR_\lambda(A)x = R_\lambda(A)Ax$, $x \in D_A$. Finally, the Yosida approximations to A are defined by

$$(3.7) \quad A_\alpha x := \frac{1}{\alpha} AR_{\frac{1}{\alpha}}(A)x, \quad x \in H.$$

Since (A, D_A) as a generator of a contractive C_0 -semigroup is m -dissipative, A_α is a special case of F_α in (3.10). The Yosida approximations A_α to A satisfy the following properties, see [10, Proposition 7.2], where

$$(3.8) \quad J_\alpha := (I - \alpha A)^{-1},$$

$J_\alpha \in B(H)$, $\|J_\alpha\| \leq 1$.

$$(3.9) \quad \begin{aligned} A_\alpha x &\xrightarrow{\alpha \rightarrow 0} Ax, \quad x \in D_A, \\ |A_\alpha x|_H &\leq |Ax|_H, \quad x \in D_A, \\ A_\alpha x &= J_\alpha Ax, \quad x \in D_A, \\ A_\alpha &\in B(H), \\ \|A_\alpha\| &\leq \frac{1}{\alpha}, \\ A_\alpha &= AJ_\alpha = \frac{1}{\alpha}(J_\alpha - I). \end{aligned}$$

Proposition 3.11. *Under Assumption 2.1*

$$\|J_\alpha\| \leq 1/(1 + \alpha\beta)$$

and

$$\langle A_\alpha x, x \rangle \leq -\beta_\alpha |x|_H^2$$

for all $x \in H$, where

$$\beta_\alpha := \frac{\beta}{1 + \alpha\beta}.$$

Proof. To prove the first inequality, let $x \in H$ and $y := J_\alpha x$, that is $x = y - \alpha Ay$. Then note that

$$|x|_H \cdot |y|_H \geq \langle x, y \rangle = \langle y - \alpha Ay, y \rangle \geq (1 + \alpha\beta)|y|_H^2,$$

which implies $|x| \geq (1 + \alpha\beta)|y|$. To prove the second inequality, note that

$$\langle -A_\alpha x, x \rangle = \langle -Ay, y - \alpha Ax \rangle \geq \beta|y|_H^2 + \alpha|Ay|_H^2 = \beta|y|_H^2 + \frac{1}{\alpha}|x - y|_H^2 \geq \frac{\beta|x|_H^2}{1 + \alpha\beta},$$

where the last inequality is obtained by minimization over all $y \in H$. \square

Note that the estimates in Proposition 3.11 are best possible under Assumption 2.1.

3.3. Yosida approximations to F . Recall that to define pseudo-weak solutions in Definition 2.5, we used the Yosida approximation to F satisfying Assumption 2.3. While there are standard references for this approximation such as [4, 5, 9]), and in the setting similar to the one considered in this paper in [15, 16, 45], we include details for completeness: fix $t \in [0, \infty)$ and set $F := F(t, \cdot)$. Then for any $\alpha > 0$ we define

$$(3.10) \quad F_\alpha := \frac{1}{\alpha}(J_\alpha(x) - x), \quad x \in H,$$

where

$$J_\alpha(x) := (I - \alpha F)^{-1}(x), \quad I(x) = x,$$

which is a nonlinear generalization of (3.8). Then each F_α is single-valued, dissipative, Lipschitz continuous with Lipschitz constant less than $\frac{2}{\alpha}$ and satisfies

$$(3.11) \quad \lim_{\alpha \rightarrow 0} F_\alpha(x) = F_0(x), \quad x \in D_F,$$

$$(3.12) \quad |F_\alpha(x)|_H \leq |F_0(x)|_H, \quad x \in D_F.$$

It is clear from the last inequality that for each $x_0 \in D_F$

$$(3.13) \quad |F_\alpha(t, x)|_H \leq |F_0(t, x_0)|_H + \frac{2}{\alpha}|x|_H \leq a(|x_0|_H) + \frac{2}{\alpha}|x|_H, \quad x \in H.$$

4. ALMOST SURE φ -TYPE ESTIMATES OF SOLUTIONS X_t

4.1. **Properties of function space \mathcal{M} .** Before proceeding to the proof of Theorem 2.8 we need to establish properties of functions in \mathcal{M} depending on the value of L_φ as introduced in Definition 2.7. In particular, we shall see that functions in \mathcal{M} satisfy the standard condition in the de la Vallée-Poussin Theorem. We also find sharp constants that might be useful for finding φ -moments depending on the growth of F_0 as measured by the radial function a in Assumption 2.4.

Lemma 4.1. *Suppose $\varphi \in \mathcal{M}$, then*

(i) *for any $c > 0$, $\beta > 0$ and any $0 < B < \beta L_\varphi$ there is a constant $C \geq 0$ such that*

$$\varphi(u) \left[\frac{\varphi'(u)}{\varphi(u)} (c\sqrt{u} - \beta u) + B \right] \leq C, \text{ for all } u \in (0, \infty).$$

The constant C can be chosen as follows.

$$(4.1) \quad C(c, \beta, B) := \max_{u \in [0, \infty)} (\varphi'(u) (c\sqrt{u} - \beta u) + B\varphi(u)) = \max_{u \in [0, u_0]} (\varphi'(u) (c\sqrt{u} - \beta u) + B\varphi(u)),$$

where $u_0 := \max \left\{ \frac{c^2}{\beta^2}, \frac{c^2}{4(\beta-B)^2} \right\}$. In particular, for $B = \frac{\beta}{2}$

$$C\left(c, \beta, \frac{\beta}{2}\right) = \frac{\beta}{2} \varphi\left(\frac{c^2}{\beta^2}\right).$$

(ii) *If $L_\varphi > 1$, then*

$$\frac{\varphi(u)}{u} \xrightarrow{u \rightarrow \infty} \infty.$$

Proof. (ii): Define $H : (0, \infty) \rightarrow (0, \infty)$ by $H(u) := \frac{\varphi(u)}{u}$. Then

$$(4.2) \quad H'(u) = \left(\frac{\varphi(u)}{u} \right)' = \left(\frac{u\varphi'(u)}{\varphi(u)} - 1 \right) \cdot \frac{\varphi(u)}{u^2} = \left(\frac{u\varphi'(u)}{\varphi(u)} - 1 \right) \cdot \frac{H(u)}{u}.$$

Using the assumption that $L_\varphi > 1$ we see that there exists a $K > 0$ such that

$$H'(u) = \left(\frac{\varphi(u)}{u} \right)' > K \frac{H(u)}{u} > 0$$

for all large enough u . Thus

$$\frac{H'(u)}{H(u)} > \frac{K}{u}$$

for such a u . Then for some $M > 0$

$$H(u) = \frac{\varphi(u)}{u} > Mu^K \text{ for all large enough } u,$$

which implies that $\frac{\varphi(u)}{u} \xrightarrow{u \rightarrow \infty} \infty$.

(i): It is enough to check that for $B \in (0, \beta L_\varphi)$

$$\varphi(u) \left[\frac{\varphi'(u)}{\varphi(u)} (c\sqrt{u} - \beta u) + B \right] \xrightarrow{u \rightarrow \infty} -\infty,$$

and so there is a $u_0 > 0$ such that

$$\varphi'(u) (c\sqrt{u} - \beta u) + B\varphi(u) < 0 \text{ for all } u > u_0.$$

Then we can choose

$$(4.3) \quad C := \max_{u \in [0, u_0]} (\varphi'(u) (c\sqrt{u} - \beta u) + B\varphi(u)) (> 0).$$

Observe that

$$\frac{\varphi'(u)}{\varphi(u)} (c\sqrt{u} - \beta u) = \frac{u\varphi'(u)}{\varphi(u)} \left(\frac{c}{\sqrt{u}} - \beta \right) \xrightarrow{u \rightarrow \infty} -\beta L_\varphi \quad (:= -\infty, \text{ if } L_\varphi = \infty),$$

and so

$$\varphi(u) \left[\frac{\varphi'(u)}{\varphi(u)} (c\sqrt{u} - \beta u) + B \right] \xrightarrow{u \rightarrow \infty} -\infty.$$

Recall that we can take C to be the maximum of the following function

$$f(u) := \varphi'(u) (c\sqrt{u} - \beta u) + B\varphi(u).$$

First we take the derivative of this function

$$\begin{aligned} f'(u) &= \varphi''(u) (c\sqrt{u} - \beta u) + \varphi'(u) \left(\frac{c}{2\sqrt{u}} - \beta \right) + B\varphi'(u) = \\ &= \varphi''(u) \sqrt{u} (c - \beta\sqrt{u}) + \varphi'(u) \left(\frac{c}{2\sqrt{u}} - (\beta - B) \right). \end{aligned}$$

By assumption φ is an increasing convex function, and therefore φ'' and φ' are non-negative, so, since $\beta - B > 0$, $f'(u) \leq 0$ for any $u \geq u_0 = \max \left\{ \frac{c^2}{\beta^2}, \frac{c^2}{4(\beta - B)^2} \right\}$. Therefore we can choose

$$C(c, \beta, B) = \max_{u \in [0, \infty)} f(u) = \max_{u \in [0, u_0]} f(u).$$

Finally, if $B = \beta/2$, then $u_0 = \frac{c^2}{\beta^2}$, and $f'(u) \geq 0$ on $[0, u_0]$, so

$$C(c, \beta, \beta/2) = f(u_0) = \frac{\beta}{2} \varphi \left(\frac{c^2}{\beta^2} \right).$$

□

We illustrate properties of functions in \mathcal{M} by considering several fundamental examples.

Example 4.2. [Polynomial functions] Suppose $\varphi(u) = u^p$, $p \geq 1$, then $\varphi \in \mathcal{M}$. In this case $L_\varphi = p$. To see how we can find C in (4.3), observe that for any $0 < B < p\beta$

$$\begin{aligned} f(u) &:= \varphi(u) \left[\frac{\varphi'(u)}{\varphi(u)} (c\sqrt{u} - \beta u) + B \right] = \\ &= cpu^{p-1/2} + (B - p\beta) u^p, \end{aligned}$$

for which

$$\begin{aligned} f'(u) &= cp \left(p - \frac{1}{2} \right) u^{p-3/2} + (B - p\beta) pu^{p-1} = \\ &= pu^{p-3/2} \left(c \left(p - \frac{1}{2} \right) - (p\beta - B) \sqrt{u} \right). \end{aligned}$$

Then the maximum of f is attained at $u_0 = \left(\frac{c(p-\frac{1}{2})}{p\beta-B} \right)^2$. Therefore

$$C_p(c, \beta, B) := \frac{c}{2} \left(\frac{c(p-\frac{1}{2})}{p\beta-B} \right)^{2p-1} = \frac{c^{2p}}{2} \left(\frac{(p-\frac{1}{2})}{p\beta-B} \right)^{2p-1}.$$

In this example $L_\varphi = p$, and so by Lemma 4.1 for any $0 < B < \beta$ we can choose

$$C_1(c, \beta, B) := \frac{c^2}{4(\beta-B)}.$$

Example 4.3. [Exponential] Suppose $\varphi(u) = e^u$, then $\varphi \in \mathcal{M}$. In this case $L_\varphi = \infty$, so we can take any positive constant B . For example, if $B = \beta/2$, then for

$$f(u) := e^u [c\sqrt{u} - \beta u + B] = e^u \left[c\sqrt{u} - \beta u + \frac{\beta}{2} \right]$$

we have

$$f'(u) = e^u \left[c\sqrt{u} + \frac{c}{2\sqrt{u}} - \beta u - \frac{\beta}{2} \right] = e^u \left(\frac{c}{\sqrt{u}} - \beta \right) \left(u + \frac{1}{2} \right)$$

and we can take

$$C = f \left(\frac{c^2}{\beta^2} \right) = \frac{\beta}{2} e^{\frac{c^2}{\beta^2}}.$$

Example 4.4. Suppose $\varphi(u) = u \ln(u+1)$, then $\varphi \in \mathcal{M}$. In this case $L_\varphi = 1$, so we can take any $0 < B < \beta$ and then C can be chosen by finding the maximum of the function

$$\begin{aligned} f(u) &:= \varphi(u) \left[\frac{\varphi'(u)}{\varphi(u)} (c\sqrt{u} - \beta u) + B \right] = \\ &\ln(u+1) (c\sqrt{u} - (\beta-B)u) + \frac{u}{u+1} (c\sqrt{u} - \beta u). \end{aligned}$$

Note that for $u > \left(\frac{c}{\beta-B} \right)^2$ the function $f(u)$ is negative. Therefore it is enough to find the maximum of f on $\left(0, \left(\frac{c}{\beta-B} \right)^2 \right)$. We will use a rough estimate for $u \in \left(0, \left(\frac{c}{\beta-B} \right)^2 \right)$

$$\begin{aligned} &\ln(u+1) (c\sqrt{u} - (\beta-B)u) + \frac{u}{u+1} (c\sqrt{u} - \beta u) \leq \\ &\frac{c^2}{4(\beta-B)} \ln(u+1) + \frac{c^2}{4\beta} \frac{u}{u+1} \leq \\ &\frac{c^2}{4(\beta-B)} u + \frac{c^2}{4\beta} \leq \frac{c^2}{4(\beta-B)} \left(\frac{c}{\beta-B} \right)^2 + \frac{c^2}{4\beta}. \end{aligned}$$

Thus we can take

$$C(c, \beta, B) := \frac{c^2}{4} \left(\frac{c^2}{(\beta-B)^3} + \frac{1}{\beta} \right).$$

4.2. Proof of Theorem 2.8. Recall now that the Ornstein-Uhlenbeck process $W_{x,A,\sigma}(t)$ defined by (2.1) is a Gaussian random variable with values in H with mean 0 and the covariance operator Q_t given by

$$Q_t x = \int_0^t e^{sA} \sigma^2 e^{sA^*} x ds.$$

We will use the following notation for the maximum process

$$(4.4) \quad W_{x,A,\sigma}^*(t) := \sup_{s \in [0,t]} |W_{x,A,\sigma}(s)|_H.$$

Proposition 4.5. *For any $t > 0$ there is an $\varepsilon > 0$ such that*

$$(4.5) \quad \mathbb{E} \left(e^{\varepsilon [W_{0,A,\sigma}^*(t)]^2} \right) < \infty.$$

Proof. This follows from Fernique's Theorem [23], see also [33, Theorem 3.1], if one can show that the law of $W_{0,A,\sigma}$ is a Gaussian measure on $C([0,t]; H)$. In the case A is self-adjoint, this follows from [34, Proposition I.0.7]. When A is not necessarily self-adjoint the same proof works using [14, Theorem 2.9]. \square

Notation 4.6. *For any $\varphi \in \mathcal{M}$ and for all $t > 0$ we denote the following random functions by*

$$K_{\varphi,\beta,a}(t) := \varphi \left(\frac{2 [a(W_{0,A,\sigma}^*(t))]^2}{\beta^2} \right),$$

$$K_\varphi(t) := \varphi \left(2 |W_{0,A,\sigma}(t)|_H^2 \right)$$

Note that these functions are finite a.s.

Remark 4.7. We will make use of the following elementary inequalities: for any $a, b \geq 0$, and $p \geq 1$

$$(4.6) \quad (a+b)^p \leq 2^{p-1} (a^p + b^p),$$

$$e^{(a+b)^2} \leq \frac{e^{4a^2}}{2} + \frac{e^{4b^2}}{2}.$$

We are now in position to prove pathwise estimates in Theorem 2.8.

Proof of Theorem 2.8. Suppose $\alpha' > 0$, $\alpha > 0$ and $Z_{\alpha',\alpha,t}^x$ is a solution to

$$(4.7) \quad \begin{aligned} dZ_{\alpha',\alpha,t}^x &= (A_{\alpha'} Z_{\alpha',\alpha,t}^x + F_\alpha(t, Z_{\alpha',\alpha,t}^x + W_{0,A,\sigma}(t))) dt, \\ Z_{\alpha',\alpha,0}^x &= x. \end{aligned}$$

Note that coefficients in (4.7) are Lipschitz, and therefore the solution exists and it is unique, and in addition the solution is continuous in t . Then for Lebesgue measure-a.e. $t > 0$ and a

C^1 function $\varphi : [0, \infty) \rightarrow (0, \infty)$

$$\begin{aligned}
 & \frac{d}{dt} \varphi (|Z_{\alpha', \alpha, t}^x|_H^2) \\
 &= 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) \langle (Z_{\alpha', \alpha, t}^x)' , Z_{\alpha', \alpha, t}^x \rangle \\
 &= 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) \langle A_{\alpha'} Z_{\alpha', \alpha, t}^x + F_{\alpha} (t, Z_{\alpha', \alpha, t}^x + W_{0, A, \sigma} (t)) , Z_{\alpha', \alpha, t}^x \rangle \\
 &= 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) \langle A_{\alpha'} Z_{\alpha', \alpha, t}^x + (F_{\alpha} (t, Z_{\alpha', \alpha, t}^x + W_{0, A, \sigma} (t)) - F_{\alpha} (t, W_{0, A, \sigma} (t))) , Z_{\alpha', \alpha, t}^x \rangle \\
 &\quad + 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) \langle F_{\alpha} (t, W_{0, A, \sigma} (t)) , Z_{\alpha', \alpha, t}^x \rangle \\
 &\leq 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) \langle A_{\alpha'} Z_{\alpha', \alpha, t}^x, Z_{\alpha', \alpha, t}^x \rangle + 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) \langle F_{\alpha} (t, W_{0, A, \sigma} (t)) , Z_{\alpha', \alpha, t}^x \rangle \\
 &\leq 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) (-\beta_{\alpha'} |Z_{\alpha', \alpha, t}^x|_H^2 + |Z_{\alpha', \alpha, t}^x|_H |F_{\alpha} (t, W_{0, A, \sigma} (t))|_H) \\
 &\leq 2\varphi' (|Z_{\alpha', \alpha, t}^x|_H^2) (a (|W_{0, A, \sigma} (t)|_H) |Z_{\alpha', \alpha, t}^x|_H - \beta_{\alpha'} |Z_{\alpha', \alpha, t}^x|_H^2) ,
 \end{aligned}$$

where we used Proposition 3.11, Equation (3.12), Assumption 2.3 and Equation (2.3). We mention that F_{α} is monotone, see for instance [12, Appendix A].

Now suppose $\varphi \in \mathcal{M}$, then by Lemma 4.1 taking $B = \beta_{\alpha'}/2$ and

$$C := C \left(a (|W_{0, A, \sigma} (t)|_H) , \beta_{\alpha'} , \frac{\beta_{\alpha'}}{2} \right) = \frac{\beta_{\alpha'}}{2} \varphi \left(\frac{[a (|W_{0, A, \sigma} (t)|_H)]^2}{\beta_{\alpha'}^2} \right) ,$$

we obtain that for all $u \in (0, \infty)$

$$\varphi' (u) (a (|W_{0, A, \sigma} (t)|_H) \sqrt{u} - \beta_{\alpha'} u) \leq C - \frac{\beta_{\alpha'}}{2} \varphi (u) ,$$

therefore

$$\frac{d}{dt} \varphi (|Z_{\alpha', \alpha, t}^x|_H^2) \leq \beta_{\alpha'} \left(\varphi \left(\frac{[a (|W_{0, A, \sigma} (t)|_H)]^2}{\beta_{\alpha'}^2} \right) - \varphi (|Z_{\alpha', \alpha, t}^x|_H^2) \right) .$$

Now by Gronwall's inequality we see that for all $t \geq 0$

$$(4.8) \quad \varphi (|Z_{\alpha', \alpha, t}^x|_H^2) \leq \varphi (|x|_H^2) e^{-\beta_{\alpha'} t} + \beta_{\alpha'} \int_0^t e^{-\beta_{\alpha'} (t-s)} \varphi \left(\frac{[a (|W_{0, A, \sigma} (s)|_H)]^2}{\beta_{\alpha'}^2} \right) ds .$$

Similarly to the proofs of [12, Lemma 1.2.3, Lemma 1.3.1 and Appendix A] one can show that $Z_{\alpha', \alpha, t}^x \rightarrow Z_{\alpha, t}^x$ locally uniformly in $t \in [0, \infty)$ for \mathbb{P} -a.e. $\omega \in \Omega$. So, since φ is continuous (4.8) holds for $Z_{\alpha, t}^x$ replacing $Z_{\alpha', \alpha, t}^x$.

Now we can use (4.6) and the fact that φ is convex to see that for the solution $X_{\alpha, t}^x$ to (2.5) we have

$$\begin{aligned}
& \varphi(|X_{\alpha,t}^x|_H^2) \\
& \leq \frac{1}{2}\varphi(2|Z_{\alpha,t}^x|_H^2) + \frac{1}{2}\varphi(2|W_{0,A,\sigma}(t)|_H^2) \\
& \leq \frac{1}{2}\varphi(2|x|_H^2) e^{-\beta t} + \frac{\beta}{2} \int_0^t e^{-\beta(t-s)} \varphi\left(\frac{2[a(|W_{0,A,\sigma}(s)|_H)]^2}{\beta^2}\right) ds + \frac{1}{2}\varphi(2|W_{0,A,\sigma}(t)|_H^2) \\
& \leq \frac{1}{2}\varphi(2|x|_H^2) e^{-\beta t} + \frac{1}{2}\varphi(2|W_{0,A,\sigma}(t)|_H^2) + \varphi\left(\frac{2[a(W_{0,A,\sigma}^*(t))]^2}{\beta^2}\right) \frac{\beta}{2} \int_0^t e^{-\beta(t-s)} ds \\
& \leq \frac{e^{-\beta t}}{2}\varphi(2|x|_H^2) + \frac{1}{2}\varphi(2|W_{0,A,\sigma}(t)|_H^2) + \frac{\beta t}{2}\varphi\left(\frac{2[a(W_{0,A,\sigma}^*(t))]^2}{\beta^2}\right).
\end{aligned}$$

Here we replaced $\varphi(\cdot)$ by $\varphi(2\cdot)$ which is again in \mathcal{M} .

Thus we have an estimate for $\varphi(|X_{\alpha,t}^x|_H^2)$ which is uniform in α , so we can apply φ^{-1} to the above inequality and use Corollary 3.9 with $(S, \mu) = ([0, \infty) \times \Omega, dt \times \mathbb{P})$ to pass to the limit as $\alpha \rightarrow 0$ along a subsequence. Then we apply φ to the resulting inequality to obtain (2.6). \square

4.3. Further *a priori* pathwise estimates of X_t and proof of Theorem 2.9. Below we prove more bounds on X_t which in particular imply Theorem 2.9. Thus we work in the setting of Theorem 2.9, and in particular we assume that Assumptions 2.1–2.4 hold.

Proposition 4.8. *Let $Z_{\alpha,t}^x$ be a solution to the regularized equation (2.4). Suppose Z_t^x is a pseudo weak limit point of $Z_{\alpha,t}^x$, $\alpha \rightarrow 0$. Then almost surely for all $\alpha > 0$*

$$(4.9) \quad |Z_{\alpha,t}^x|_H \leq |x|_H e^{-\beta t} + \frac{1}{2} \int_0^t e^{-\beta(t-s)} a(|W_{0,A,\sigma}(s)|_H) ds =: Z_t^{*,x}$$

for all $t \geq 0$ and thus

$$(4.10) \quad |Z^x|_H \leq Z^{*,x} \quad (dt \times \mathbb{P})\text{-a.s.}$$

Proof. One of the observations in the proof of Theorem 2.8 was that (4.8) holds for $Z_{\alpha',\alpha,t}^x$ instead of $Z_{\alpha,t}^x$. Now we can take φ to be the identity map and apply [19, Theorem 5] to obtain (4.9). Equation (4.10) then follows by Corollary 3.9 with $(S, \mu) = ([0, \infty) \times \Omega, dt \times \mathbb{P})$. \square

Proof of Theorem 2.9. The assertion follows from (4.9) and Proposition 3.10 with $(S, \mu) = ([0, \infty) \times \Omega, dt \times \mathbb{P})$. \square

5. UNIFORM INTEGRABILITY OF GIRSANOV DENSITIES

5.1. Stopping times and Girsanov transforms. Fix $T > 0$ and define a sequence of stopping times by

$$(5.1) \quad \tau_n^x := \inf\{t \in [0, T] : Z_t^{*,x} + |W_{0,A,\sigma}(t)|_H > n\} \wedge T, n \in \mathbb{N}.$$

Note that stopping times τ_n^x do not depend on α and that \mathbb{P} -almost surely

$$(5.2) \quad \lim_{n \rightarrow \infty} \tau_n^x = T.$$

Note also that $Z_t^{*,x} + |W_{0,A,\sigma}(t)|_H$ as a \mathbb{P} -a.s. continuous process.

The following lemma is used in the proof of Theorem 2.10 in Section 5.2.

Lemma 5.1. *We have that for any $n > 3|x|_H$*

$$\mathbb{1}_{\{t \in [0, \tau_n^x]\}} |X_{t, \alpha}^x|_H \leq n \quad \text{for all } \alpha > 0, t \geq 0.$$

Proof. This follows immediately from (4.9). \square

Now we consider Girsanov transforms for the Yosida regularized equations (2.4). For $x \in H$ let

$$(5.3) \quad \rho_\alpha(x, t) := \exp(\zeta_\alpha(x, t)),$$

where

$$(5.4) \quad \begin{aligned} \zeta_\alpha(x, t) := & \int_0^t \langle \sigma^{-1} F_\alpha(s, W_{x, A, \sigma}(s)), dW(s) \rangle \\ & - \frac{1}{2} \int_0^t |\sigma^{-1} F_\alpha(s, W_{x, A, \sigma}(s))|_H^2 ds. \end{aligned}$$

We define the measure \mathbb{P}_α^x on $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$ by

$$(5.5) \quad \frac{d\mathbb{P}_\alpha^x}{d\mathbb{P}} = \rho_\alpha(x, T) =: \rho_\alpha^x,$$

and we denote by \mathbb{E}_α^x the expectation with respect to the probability measure \mathbb{P}_α^x given by (5.5). Note that this gives a weak mild solution to (2.5) according to [34, Appendix I]. More precisely, we define

$$(5.6) \quad \tilde{X}_x(t) := W_{x, A, \sigma}(t).$$

Note that this process does not depend on α although the measure \mathbb{P}_α^x does depend on α which is important in (5.8) below. Denote

$$(5.7) \quad \tilde{W}_{x, \alpha}(t) := W_t - \int_0^t \sigma^{-1} F_\alpha(s, W_{x, A, \sigma}(s)) ds,$$

then $\tilde{W}_{x, \alpha}$ is a cylindrical Wiener process under \mathbb{P}_α^x and

$$(5.8) \quad \begin{aligned} d\tilde{X}_x(t) &= dW_{x, A, \sigma}(t) = A\tilde{X}_x dt + \sigma dW_t \\ &= A\tilde{X}_x dt + F_\alpha(t, \tilde{X}_x(t)) dt + \sigma d\tilde{W}_{x, \alpha}(t), \end{aligned}$$

that is, $\tilde{X}_x(t)$ solves this equation in the mild sense.

Remark 5.2 (On localization). As a side remark we would like to mention that in infinite dimensions the processes defined by (5.8) are not semimartingales in general (unlike in [36]). One might want to use localization to introduce

$$(5.9) \quad \tilde{W}_{x, \alpha}^n(t) = W_t - \int_0^{t \wedge \tau_n} \sigma^{-1} F_\alpha(s, W_{x, A, \sigma}(s)) ds,$$

$$(5.10) \quad \rho_\alpha^n(x, t) = \exp(\zeta_\alpha(x, t \wedge \tau_n^x)),$$

and then define $\rho_\alpha(x, t)$ as a limit as $n \rightarrow \infty$, if the limit exists. However, the localization can not be used easily for the equations with non-smooth coefficients because interchanging the limits as $n \rightarrow \infty$ and $\alpha \rightarrow 0$ may be problematic. We use stopping times in a different way in (5.17).

5.2. Estimates of the Girsanov densities and proof of Theorem 2.10.

Proof of Theorem 2.10. In this proof we assume that $x \in D_F$ and $T > 0$ are fixed, and $t \in [0, T]$. Subsequently we abuse notation, and drop dependence on x, T , although our estimates depend on x and T .

By (5.5) we have that for any Borel-measurable $\Psi : [0, \infty) \rightarrow [0, \infty)$

$$(5.11) \quad \mathbb{E} \rho_\alpha^x \Psi(\rho_\alpha^x) = \mathbb{E}_\alpha^x \Psi(\rho_\alpha^x),$$

where ρ_α^x is the density defined by (5.5). Note that by [37] the distribution of $(\widetilde{W}_{x,\alpha}, \widetilde{X}_x)$ under the measure \mathbb{P}_α^x is the same as the distribution of (W, X_α^x) under the measure \mathbb{P} as we pointed out in Section 5.1. In particular, by Assumption 2.3

$$(5.12) \quad X_\alpha^x \in D_F \quad dt \times \mathbb{P} - \text{a.s.}$$

Recall that $(\widetilde{W}_{x,\alpha}, \widetilde{X}_x) = (\widetilde{W}_{x,\alpha}, W_{x,A,\sigma})$. Then

$$(5.13) \quad \mathbb{E}_\alpha^x \Psi(\rho_\alpha^x) = \mathbb{E} \Psi(\widetilde{\rho}_\alpha^x),$$

where

$$(5.14) \quad \widetilde{\rho}_\alpha^x := \exp \left(\int_0^t \langle \sigma^{-1} F_\alpha(s, X_\alpha^x(s)), dW(s) \rangle + \frac{1}{2} \int_0^t |\sigma^{-1} F_\alpha(s, X_\alpha^x(s))|_H^2 ds \right).$$

We can estimate $\mathbb{E} \mathbb{1}_A |\widetilde{\rho}_\alpha^x|^p$ for any $A \in \mathcal{F}$ as follows.

$$\begin{aligned} & \mathbb{E} \mathbb{1}_A |\widetilde{\rho}_\alpha^x|^p = \\ & \mathbb{E} \mathbb{1}_A \exp \left(p \int_0^t \langle \sigma^{-1} F_\alpha(s, X_\alpha^x(s)), dW_s \rangle - p^2 \int_0^t |\sigma^{-1} F_\alpha(s, X_\alpha^x(s))|_H^2 ds \right) \\ & \times \exp \left(\left(p^2 + \frac{p}{2} \right) \int_0^t |\sigma^{-1} F_\alpha(s, X_\alpha^x(s))|_H^2 ds \right) \\ & \leq \left(\mathbb{E} \exp \left(2p \int_0^t \langle \sigma^{-1} F_\alpha(s, X_\alpha^x(s)), dW_s \rangle - 2p^2 \int_0^t |\sigma^{-1} F_\alpha(s, X_\alpha^x(s))|_H^2 ds \right) \right)^{1/2} \\ & \times \left(\mathbb{E} \mathbb{1}_A \exp \left((2p^2 + p) \int_0^t |\sigma^{-1} F_\alpha(s, X_\alpha^x(s))|_H^2 ds \right) \right)^{1/2}. \end{aligned}$$

Note that the first term in the last equation is equal to the expectation of the stochastic exponential for the martingale $2p \int_0^t \langle \sigma^{-1} F_\alpha(s, X_\alpha^x(s)), dW_s \rangle$, and so its expectation is 1. Therefore,

$$(5.15) \quad \mathbb{E} \mathbb{1}_A |\widetilde{\rho}_\alpha^x|^p \leq \left(\mathbb{E} \mathbb{1}_A \exp \left((2p^2 + p) \int_0^t |\sigma^{-1} F_\alpha(s, X_\alpha^x(s))|_H^2 ds \right) \right)^{1/2}.$$

Let us first assume that $\sup_{r>0} a(r) < \infty$. Then taking $A := \{\tau_n^x \geq t\}$, $n > 3|x|_H$, it follows by (5.12), (3.9) and Assumption 2.4 that the right-hand side in (5.15) is bounded uniformly in α , and therefore we can choose $\Psi(y) = y^p$ for any $p > 0$ to conclude by (5.11) that ρ_α^x , $\alpha > 0$, are uniformly $L^1(\Omega, \mathbb{P})$ -integrable. Now let us assume that

$$\sup_{r>0} a(r) = \infty.$$

Then we can construct Ψ as follows. We take $p = 2$ and use Assumption 2.4, Equations (5.11), (5.12), (5.13), Proposition 4.8 and Lemma 5.1, to deduce that

$$(5.16) \quad \mathbb{E} |\widetilde{\rho}_\alpha^x|^2 \mathbb{1}_{\{\tau_n^x \geq t\}} \leq \exp\left(5(a(n)\|\sigma^{-1}\|)^2 T\right).$$

By (5.15) and Chebyshev's inequality we have

$$(5.17) \quad \begin{aligned} \mathbb{P}(\widetilde{\rho}_\alpha^x > y) &\leq \mathbb{P}(\widetilde{\rho}_\alpha^x > y, \tau_n^x \geq t) + \mathbb{P}(\tau_n^x < t) \\ &\leq \exp\left(5(a(n)\|\sigma^{-1}\|)^2 T\right) / y^2 + \mathbb{P}(\tau_n^x < T). \end{aligned}$$

For $y \in (1 + \exp(5(a(0)\|\sigma^{-1}\|)^2 T), \infty)$ we define $n(y)$ as the maximal natural number such that

$$\exp\left(5(a(n(y))\|\sigma^{-1}\|)^2 T\right) < y$$

or

$$(5.18) \quad a(n(y)) < \frac{1}{\|\sigma^{-1}\|} \left(\frac{\log(y)}{5T}\right)^{1/2}.$$

Recall that we assumed in this part of the proof that $a(n)$ is non-decreasing and unbounded, and therefore we have that $n(y)$ is non-decreasing in y and $\lim_{y \rightarrow \infty} n(y) = \infty$. Then we can define the non-increasing function by

$$(5.19) \quad p_0(y) := p_0(y, \sigma, A, a, x, T) := \min\{1, 1/y + \mathbb{P}(\tau_{n(y)}^x < T)\}, \quad y \in (0, \infty).$$

Observe that by (5.2)

$$\lim_{y \rightarrow \infty} p_0(y, \sigma, A, a, x, T) = 0.$$

Suppose $\Psi : [0, \infty) \rightarrow [0, \infty)$ is an increasing function which is C^1 on $(0, \infty)$ with $\Psi(0) = 0$. Then obviously

$$(5.20) \quad \mathbb{E}\Psi(Y) = \mathbb{E} \int_0^\infty \Psi'(y) \mathbb{1}_{\{0 < y < Y\}} dy = \int_0^\infty \Psi'(y) \mathbb{P}\{Y > y\} dy$$

for any non-negative random variable Y . We apply this equation to $Y = \widetilde{\rho}_\alpha^x$ to see that by (5.17), (5.19) for all $\alpha > 0$

$$(5.21) \quad \mathbb{E}\Psi(\widetilde{\rho}_\alpha^x) \leq \int_0^\infty \Psi'(y) p_0(y) dy.$$

Thus by (5.11) and (5.13) this implies uniform $L^1(\Omega, \mathbb{P})$ -integrability of ρ_α^x , $\alpha > 0$, if we can find Ψ as above with the following two properties

$$(5.22) \quad \int_0^\infty \Psi'(y) p_0(y) dy < \infty$$

and

$$(5.23) \quad \lim_{y \rightarrow \infty} \Psi(y) = \infty.$$

The existence of such a Ψ can be seen as follows: since $y \mapsto p_0(y)$ decreases to zero as $y \rightarrow \infty$, we can find a sequence $\{y_k\}_{k \in \mathbb{N}}$ in $(0, \infty)$ such that

$$y_k + 3 < y_{k+1}, \quad k \in \mathbb{N},$$

and $p_0(y) \leq \frac{1}{k^2}$ for $y \geq y_k$. Now define $g : [0, \infty) \rightarrow [0, \infty)$ by

$$g = \sum_{k=1}^{\infty} \Psi_k,$$

where $\Psi_k \in C^\infty((0, \infty))$ such that

$$\mathbb{1}_{[y_k+1, y_k+2]} \leq \Psi_k \leq \mathbb{1}_{[y_k, y_k+3]}.$$

Define

$$\Psi(y) = \int_0^y g(s) ds, \quad y \geq 0.$$

Then $\Psi : [0, \infty) \rightarrow [0, \infty)$ is continuous, increasing and C^∞ on $(0, \infty)$. Furthermore, obviously (5.22) and (5.23) hold. Another construction of a function Ψ is given in the beginning of Subsection 5.3.

Now let $X^x \in L^2(\Omega, \mathbb{P}; L^2([0, T], dt; H)) = L^2([0, T] \times \Omega, dt \times \mathbb{P}; H)$ be the pseudo-weak limit of $X_{\alpha_n}^x$ as $\alpha_n \xrightarrow{n \rightarrow \infty} 0$, with the corresponding function $\psi : H \rightarrow H$ defined by (3.2). By the above and the Dunford-Pettis theorem, choosing another subsequence if necessary we have

$$\rho_{\alpha_n}^x \xrightarrow{n \rightarrow \infty} \rho^x \text{ in } L^1(\Omega, \mathbb{P})$$

for some $\rho^x \in L^1(\Omega, \mathbb{P})$.

Let $\mathcal{X} := L^2([0, T]; H)$ and let $G : \mathcal{X} \rightarrow \mathbb{R}$ be bounded and sequentially weakly continuous. Then for $\mathbb{Q}_x := \mathbb{P} \circ W_{x, A, \sigma}^{-1}$

$$\begin{aligned} \int_{\mathcal{X}} G \circ \psi d(\mathbb{P} \circ (X^x)^{-1}) &= \int_{\Omega} G(\psi(X^x)) d\mathbb{P} = \lim_{n \rightarrow \infty} \int_{\Omega} G(\psi(X_{\alpha_n}^x)) d\mathbb{P} \\ (5.24) \qquad \qquad \qquad &= \lim_{n \rightarrow \infty} \int_{\Omega} (G \circ \psi)(W_{x, A, \sigma}) \rho_{\alpha_n}^x d\mathbb{P} = \int_{\Omega} (G \circ \psi)(W_{x, A, \sigma}) \rho^x d\mathbb{P} \\ &= \int_{\Omega} (G \circ \psi)(W_{x, A, \sigma}) \mathbb{E}_{\mathbb{P}}[\rho^x | W_{x, A, \sigma}] d\mathbb{P} = \int_{\mathcal{X}} G \circ \psi \bar{\rho}^x d\mathbb{Q}_x, \end{aligned}$$

where

$$\bar{\rho}^x := \mathbb{E}_{\mathbb{P}}[\rho^x | W_{x, A, \sigma} = \cdot].$$

Let \mathcal{N} denote the set of all such functions $G \circ \psi : H \rightarrow \mathbb{R}$ from above. Since ψ in (3.2) is one-to-one, we can find a countable set $\mathcal{N}_0 \subset \mathcal{N}$, which separates the points in \mathcal{X} . Indeed, let $\{e_i\}_{i \in \mathbb{N}}$ and $\{g_i\}_{i \in \mathbb{N}}$ be orthonormal bases of H and $L^2([0, T], dt; \mathbb{R})$ respectively. Define maps $G_{ij} : \mathcal{X} \rightarrow \mathbb{R}$, $i, j \in \mathbb{N}$,

$$G_{ij}(w) := \int_0^T g_i(t) \langle e_j, w(t) \rangle_H dt, \quad w \in \mathcal{X}.$$

Then obviously $\{G_{ij}, i, j \in \mathbb{N}\}$ separates the points of \mathcal{X} and hence so does $\mathcal{N}_0 := \{(N \wedge G_{ij} \vee (-N)) \circ \psi, i, j, N \in \mathbb{N}\}$. Clearly, each $N \wedge G_{ij} \vee (-N)$ is weakly continuous, so $\mathcal{N}_0 \subset \mathcal{N}$. Furthermore, obviously \mathcal{N} is closed under multiplication and consists of bounded Borel measurable functions on \mathcal{X} . Therefore, (5.24) implies that

$$\mathbb{P} \circ (X^x)^{-1} = \bar{\rho}^x \mathbb{Q}_x.$$

Note that by Kuratowski's Theorem (e.g. [38, Section I.3]) $C([0, T]; H)$ is a Borel subset of \mathcal{X} such that $\mathbb{Q}_x(C([0, T]; H)) = 1$, therefore

$$(\mathbb{P} \circ (X^x)^{-1})(C([0, T]; H)) = 1,$$

so X^x has continuous sample paths \mathbb{P} -a.s. □

5.3. Quantitative estimates and examples. As in Subsection 5.2 we fix $x \in D_F$ and $T > 0$ below, and omit dependence on x, T in the notation.

One of the consequences of the proof of Equation (2.9) in Theorem 2.10 is a constructive approach to finding a function $\Psi(\cdot)$, though it seems not possible to find an optimal form of the function $\Psi(\cdot)$ under conditions of Theorem 2.10. Below we present some explicit results. In particular, we explain how to treat the case when the growth of the nonlinearity given by function $a(\cdot)$ is polynomial.

We begin with the following simple observations about real-valued functions. Assume that we are given a non-increasing function $p_0 : [0, \infty) \rightarrow (0, \infty)$ which is continuous at zero and such that

$$(5.25) \quad \lim_{y \rightarrow \infty} p_0(y) = 0.$$

Our aim is to find a non-decreasing absolutely continuous function $\Psi : [0, \infty) \rightarrow [0, \infty)$ such that (5.22) and (5.23) hold. If $p_0(\cdot)$ is absolutely continuous, then we can simply choose an absolutely continuous $\Psi : [0, \infty) \rightarrow [0, \infty)$ satisfying

$$(5.26) \quad \Psi(y) \leq (p_0(y))^{-1/2}$$

and show that

$$(5.27) \quad \int_0^\infty \Psi'(y)p_0(y)dy = -\Psi(0)p_0(0) - \int_0^\infty \Psi(y)p_0'(y)dy \leq 2\sqrt{p_0(0)} - \Psi(0)p_0(0).$$

Even if $p_0(\cdot)$ is not absolutely continuous, we can define for $y \in \mathbb{R}$ and $\delta \in (0, \infty)$

$$(5.28) \quad p_\delta(y) := \frac{1}{\delta} \int_{\mathbb{R}} p_0(s - \delta) m\left(\frac{s - y}{\delta}\right) ds \geq p_0(y),$$

where $m(\cdot)$ is a standard mollifier on \mathbb{R} , that is, a smooth non-negative function $m : \mathbb{R} \rightarrow [0, \infty)$ with support in $[-1, 1]$ and L^1 -norm equal to 1. We assume that in this formula $p_0(y) = p_0(0)$ if $y \leq 0$. Equation (5.28) is different from the usual mollification because of the shift by δ in $p_0(s - \delta)$, which ensures that $p_\delta(y)$ is a non-decreasing function of δ . Then $p_\delta(y)$ satisfies (5.25), and this implies the existence of a smooth function $\Psi(y) = (p_\delta(y))^{-1/2}$ satisfying (5.26).

Remark 5.3. For any fixed σ, A, a, x, T one can estimate $\Psi(y)$ in (2.9). However, in the most general form this computation is cumbersome and therefore it is not presented in our paper. Instead we illustrate this approach by giving several examples satisfying natural extra assumptions.

The following is a corollary from the proof of Theorem 2.10 given in Subsection 5.2.

Example 5.4. *Suppose*

$$(5.29) \quad a(y) \leq c(1 + y)^{n+1}$$

for some $c \geq 1, n > 0$ and all $y \geq 0$. Then we can choose $\delta_0 > 0$ such that

$$(5.30) \quad \Psi(y) := \exp\left(\left(\log(1+y)\right)^{\delta_0}\right)$$

satisfies (2.9).

Proof. Our aim is to show that for $\delta_0 > 0$ small enough, if we choose $\Psi(y)$ as in (5.30), it satisfies (5.26) and (5.27).

We define a smooth strictly increasing function $a_0 : [0, \infty) \rightarrow [0, \infty)$ by

$$(5.31) \quad a_0(y) := 1 + y + \int_{-1}^{\infty} a(s+1) m(s-y) ds > a(y) + y,$$

where $m(\cdot)$ is a standard mollifier on \mathbb{R} with support in $[-1, 1]$.

Then for $y \in (\exp(5(a_0(0)\|\sigma^{-1}\|)^2 T), \infty)$ we define

$$(5.32) \quad n(y) := \left[a_0^{-1} \left(\frac{1}{\|\sigma^{-1}\|} \left(\frac{\log(y)}{5T} \right)^{1/2} \right) \right],$$

where $[z]$ is the integer part of z and $a_0^{-1}(\cdot)$ is the inverse function of the function $a_0(\cdot)$. Note that

$$(5.33) \quad \exp\left(5(a(n(y))\|\sigma^{-1}\|)^2 T\right) < \exp\left(5(a_0(n(y))\|\sigma^{-1}\|)^2 T\right) \leq y$$

and

$$(5.34) \quad \exp\left(5(a_0(n(y)+1)\|\sigma^{-1}\|)^2 T\right) > y$$

for all $y \in (\exp(5(a_0(0)\|\sigma^{-1}\|)^2 T), \infty)$. Equation (5.29) implies that

$$a_0(y) \leq c_0(1+y)^{n+1}$$

for some $c_0 \geq c$ and all $y \in (1 + \exp(5(a_0(0)\|\sigma^{-1}\|)^2 T), \infty)$.

Recall that $W_{0,A,\sigma}^* = \sup_{t \in [0,T]} |W_{0,A,\sigma}(t)|_H$. Applying Markov's inequality to Fernique's Theorem as formulated in (4.5), we have that

$$\mathbb{P}(W_{0,A,\sigma}^* > s) < \exp(-\varepsilon s^2) \mathbb{E}\left(e^{\varepsilon [W_{0,A,\sigma}^*(T)]^2}\right)$$

for all $s > 0$. If $p_0(y)$ is given by (5.19) then, by (5.26) and (5.27), it is enough to verify the following inequality

$$p_0(y) \leq 2\mathbb{P}\left(W_{0,A,\sigma}^* \geq C_1 a_0^{-1}\left(a_0^{-1}\left(\sqrt{\log(2+y)}\right)\right)\right)$$

for some constant $C_1 > 0$ and all $y \in (y_0, \infty)$ for some $y_0 > 0$. It is enough to show that

$$\mathbb{P}(\tau_{n(y)}^x < T) \leq \mathbb{P}\left(W_{0,A,\sigma}^* \geq C_1 a_0^{-1}\left(a_0^{-1}\left(\sqrt{\log(2+y)}\right)\right)\right).$$

By definition of stopping times in (5.1) the left-hand side is the same as

$$\mathbb{P}\left(\sup_{t \in [0,T]} Z_t^{*,x} + W_{0,A,\sigma}^* > n(y)\right),$$

which, by (5.31), (5.32), (5.33), (5.34) is controlled up to constants by

$$\mathbb{P}\left(\sup_{t \in [0,T]} Z_t^{*,x} + W_{0,A,\sigma}^* > a_0^{-1}\left(\frac{1}{\|\sigma^{-1}\|} \left(\frac{\log(y)}{5T}\right)^{1/2}\right)\right).$$

Using (4.9) we can find positive constants D_1, D_2 such that this probability is smaller than

$$\begin{aligned} & \mathbb{P} \left(a_0 (D_1 + D_2 W_{0,A,\sigma}^*) \geq a_0^{-1} \left(\frac{1}{\|\sigma^{-1}\|} \left(\frac{\log(y)}{5T} \right)^{1/2} \right) \right) = \\ & \mathbb{P} \left(W_{0,A,\sigma}^* \geq \frac{1}{D_2} a_0^{-1} \left(a_0^{-1} \left(\frac{1}{\|\sigma^{-1}\|} \left(\frac{\log(y)}{5T} \right)^{1/2} \right) - \frac{D_1}{D_2} \right) \right). \end{aligned}$$

This proves (5.30) for a large enough constant D_2 and small enough constant δ_0 . \square

In particular, this estimate applies for the example

$$(5.35) \quad F(t, x) = -x|x|_H^n.$$

More generally this estimate applies to any function

$$(5.36) \quad F(t, x) = -xf(t, |x|_H)$$

where $f(t, y) \geq 0$ is a real-valued function such that for any $t \geq 0, y \geq 0$ we have $f(t, y) < c(1+y)^n$. Note that F is monotone if for any fixed $t \geq 0$ function $f(t, y)$ is increasing lower semicontinuous for $y \geq 0$. This follows from the sub-gradient representation

$$(5.37) \quad F(t, x) = -\partial \int_0^{|x|_H} f(t, s) ds,$$

see [41, §5 and §24] and [5, Section 1.2].

Remark 5.5. With more tedious computations, which we do not present in our paper, in more general situations Ψ can be defined as

$$(5.38) \quad \Psi(y) := \left(\mathbb{P} \left(W_{0,A,\sigma}^* \geq C_1 a_0^{-1} \left(C_2 a_0^{-1} \left(\sqrt{C_3 \log(2+y)} \right) \right) \right) \right)^{\varepsilon_1 - 1}.$$

for some constants $C_1, C_2, C_3 > 0$ and $\varepsilon_1 > 0$.

REFERENCES

1. S. Albeverio and Michael Röckner, *Dirichlet forms, quantum fields and stochastic quantisation*, Tech. report, BiBoS, 1988.
2. Sergio Albeverio and Raphael Høegh-Krohn, *Some remarks on Dirichlet forms and their applications to quantum mechanics and statistical mechanics*, Functional analysis in Markov processes (Katata/Kyoto, 1981), Lecture Notes in Math., vol. 923, Springer, Berlin-New York, 1982, pp. 120–132. MR 661620
3. Sergio Albeverio and Michael Röckner, *Classical Dirichlet forms on topological vector spaces—closability and a Cameron-Martin formula*, J. Funct. Anal. **88** (1990), no. 2, 395–436. MR 1038449
4. Viorel Barbu, *Nonlinear semigroups and differential equations in Banach spaces*, Editura Academiei Republicii Socialiste România, Bucharest, 1976, Translated from the Romanian. MR 0390843 (52 #11666)
5. ———, *Nonlinear differential equations of monotone types in Banach spaces*, Springer Monographs in Mathematics, Springer, New York, 2010. MR 2582280 (2011d:34001)
6. Fabrice Baudoin, Qi Feng, and Maria Gordina, *Integration by parts and quasi-invariance for the horizontal Wiener measure on foliated compact manifolds*, Journal of Functional Analysis **277** (2019), no. 5, 1362 – 1422.
7. Fabrice Baudoin, Maria Gordina, and Tai Melcher, *Quasi-invariance for heat kernel measures on sub-Riemannian infinite-dimensional Heisenberg groups*, Trans. Amer. Math. Soc. **365** (2013), no. 8, 4313–4350. MR 3055697
8. Vladimir I. Bogachev, Giuseppe Da Prato, and Michael Röckner, *Regularity of invariant measures for a class of perturbed Ornstein-Uhlenbeck operators*, NoDEA Nonlinear Differential Equations Appl. **3** (1996), no. 2, 261–268. MR 1385887 (97c:60189)

9. H. Brézis, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973, North-Holland Mathematics Studies, No. 5. Notas de Matemática (50). MR 0348562
10. Haim Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Universitext, Springer, New York, 2011. MR 2759829
11. Felix E. Browder, *Nonlinear accretive operators in Banach spaces*, Bull. Amer. Math. Soc. **73** (1967), 470–476. MR 0212626
12. Sandra Cerrai, *Second order PDE's in finite and infinite dimension*, Lecture Notes in Mathematics, vol. 1762, Springer-Verlag, Berlin, 2001, A probabilistic approach. MR MR1840644 (2002j:35327)
13. G. Da Prato, S. Kwapien, and J. Zabczyk, *Regularity of solutions of linear stochastic equations in Hilbert spaces*, Stochastics **23** (1987), no. 1, 1–23. MR 920798
14. Giuseppe Da Prato, *Kolmogorov equations for stochastic PDEs*, Advanced Courses in Mathematics. CRM Barcelona, Birkhäuser Verlag, Basel, 2004. MR 2111320 (2005m:60002)
15. Giuseppe Da Prato and Michael Röckner, *Singular dissipative stochastic equations in Hilbert spaces*, Probab. Theory Related Fields **124** (2002), no. 2, 261–303. MR 1936019 (2003k:60151)
16. Giuseppe Da Prato, Michael Röckner, and Feng-Yu Wang, *Singular stochastic equations on Hilbert spaces: Harnack inequalities for their transition semigroups*, J. Funct. Anal. **257** (2009), no. 4, 992–1017. MR 2535460 (2010k:60211)
17. Giuseppe Da Prato and Jerzy Zabczyk, *Nonexplosion, boundedness, and ergodicity for stochastic semilinear equations*, J. Differential Equations **98** (1992), no. 1, 181–195. MR 1168978
18. ———, *Stochastic equations in infinite dimensions*, second ed., Encyclopedia of Mathematics and its Applications, vol. 152, Cambridge University Press, Cambridge, 2014. MR 3236753
19. Sever Silvestru Dragomir, *Some Gronwall type inequalities and applications*, Nova Science Publishers, Inc., Hauppauge, NY, 2003. MR 2016992
20. Bruce K. Driver and Maria Gordina, *Heat kernel analysis on infinite-dimensional Heisenberg groups*, J. Funct. Anal. **255** (2008), no. 9, 2395–2461. MR MR2473262
21. ———, *Integrated Harnack inequalities on Lie groups*, J. Differential Geom. **83** (2009), no. 3, 501–550. MR MR2581356
22. Klaus-Jochen Engel and Rainer Nagel, *One-parameter semigroups for linear evolution equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000, With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt. MR 1721989 (2000i:47075)
23. X. Fernique, *Régularité des trajectoires des fonctions aléatoires gaussiennes*, Lecture Notes in Math. (1975), 1–96. Lecture Notes in Math., Vol. 480. MR 0413238
24. Dariusz Gątarek and Benjamin Goldys, *On solving stochastic evolution equations by the change of drift with application to optimal control*, Stochastic partial differential equations and applications (Trento, 1990), Pitman Res. Notes Math. Ser., vol. 268, Longman Sci. Tech., Harlow, 1992, pp. 180–190. MR 1222696 (94d:60096)
25. ———, *On invariant measures for diffusions on Banach spaces*, Potential Anal. **7** (1997), no. 2, 539–553. MR 1467205 (98k:60102)
26. Maria Gordina, *An application of a functional inequality to quasi-invariance in infinite dimensions*, pp. 251–266, Springer New York, New York, NY, 2017.
27. Maria Gordina, Michael Röckner, and Feng-Yu Wang, *Dimension-independent Harnack inequalities for subordinated semigroups*, Potential Anal. **34** (2011), no. 3, 293–307. MR 2782975 (2012f:60230)
28. Tosio Kato, *Nonlinear semigroups and evolution equations*, J. Math. Soc. Japan **19** (1967), 508–520. MR 0226230
29. Yukio Kōmura, *Nonlinear semi-groups in Hilbert space*, J. Math. Soc. Japan **19** (1967), 493–507. MR 0216342
30. N. V. Krylov, *A simple proof of the existence of a solution to the Itô equation with monotone coefficients*, Teor. Veroyatnost. i Primenen. **35** (1990), no. 3, 576–580. MR 1091217
31. ———, *Introduction to the theory of diffusion processes*, Translations of Mathematical Monographs, vol. 142, American Mathematical Society, Providence, RI, 1995, Translated from the Russian manuscript by Valim Khidkel and Gennady Pasechnik. MR 1311478
32. ———, *A simple proof of a result of A. Novikov*, an arxiv preprint, 2002.

33. Hui Hsiung Kuo, *Gaussian measures in Banach spaces*, Springer-Verlag, Berlin, 1975, Lecture Notes in Mathematics, Vol. 463. MR MR0461643 (57 #1628)
34. Wei Liu and Michael Röckner, *Stochastic partial differential equations: an introduction*, Universitext, Springer, Cham, 2015. MR 3410409
35. Tai Melcher, *Heat kernel analysis on semi-infinite Lie groups*, J. Funct. Anal. **257** (2009), no. 11, 3552–3592. MR 2572261 (2011b:58074)
36. Michel Métivier, *Semimartingales*, de Gruyter Studies in Mathematics, vol. 2, Walter de Gruyter & Co., Berlin, 1982, A course on stochastic processes. MR MR688144 (84i:60002)
37. Martin Ondreját, *Uniqueness for stochastic evolution equations in Banach spaces*, Dissertationes Math. (Rozprawy Mat.) **426** (2004), 63. MR 2067962 (2005e:60133)
38. K. R. Parthasarathy, *Probability measures on metric spaces*, AMS Chelsea Publishing, Providence, RI, 2005, Reprint of the 1967 original. MR 2169627
39. A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, Applied Mathematical Sciences, vol. 44, Springer-Verlag, New York, 1983. MR MR710486 (85g:47061)
40. R. T. Rockafellar, *On the maximality of sums of nonlinear monotone operators*, Trans. Amer. Math. Soc. **149** (1970), 75–88. MR 0282272
41. R. Tyrrell Rockafellar, *Convex analysis*, Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J., 1970. MR 0274683
42. Michael Röckner and Feng-Yu Wang, *General extinction results for stochastic partial differential equations and applications*, J. Lond. Math. Soc. (2) **87** (2013), no. 2, 545–560. MR 3046285
43. Wilhelm Stannat, *(Nonsymmetric) Dirichlet operators on L^1 : existence, uniqueness and associated Markov processes*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **28** (1999), no. 1, 99–140. MR 1679079
44. ———, *The theory of generalized Dirichlet forms and its applications in analysis and stochastics*, Mem. Amer. Math. Soc. **142** (1999), no. 678, viii+101. MR 1632609
45. Sven Wiesinger, *Uniqueness of solutions to Fokker-Planck equations related to singular SPDE driven by Lévy and cylindrical Wiener noise*, Ph.D. thesis, University of Bielefeld, 2011.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269, U.S.A.
Email address: maria.gordina@uconn.edu

‡ DEPARTMENT OF MATHEMATICS, BIELEFELD UNIVERSITY, D-33501 BIELEFELD, GERMANY
Email address: roeckner@math.uni-bielefeld.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269, U.S.A.
Email address: teplyaev@uconn.edu